

Synthesis and Crystal Structure of Nb_{0.84}N

Nancy Frenzel^a, Elisabeth Irran^a, Martin Lerch^a, and Alexandra Buchsteiner^b

^a Institut für Chemie, Technische Universität Berlin, Straße des 17. Juni 135, 10623 Berlin, Germany

^b Helmholtz-Zentrum Berlin für Materialien und Energie, Hahn-Meitner-Platz 1, 14109 Berlin, Germany

Reprint requests to Dr. Elisabeth Irran. Fax: +49 30 314 79656. E-mail: elisabeth.irran@tu-berlin.de

Z. Naturforsch. **2011**, *66b*, 1 – 6; received August 11, 2010

A new compound of the composition Nb_{0.84}N was prepared by ammonolysis of NbO₂ at 1100 °C. The crystal structure refinement was performed by the Rietveld method using X-ray and neutron powder diffraction data. Nb_{0.84}N crystallizes in the trigonal space group $R\bar{3}m$ (no. 166) with the lattice parameters $a = 298.5(2)$ and $c = 2384.3(4)$ pm. The niobium atoms form a close packing with a layer sequence which can be described by the Jagodzinski symbol hhc . The nitrogen atoms fill all octahedral voids. Along [001] a sequence of two layers of trigonal NbN₆ prisms and one layer of NbN₆ octahedra is formed. The nitrogen positions are fully occupied, the niobium positions only partially. Nb_{0.84}N is part of a family of crystal structures between the *anti*-NiAs and the NaCl type consisting of close-packed metal layers with varying stacking sequences.

Key words: Niobium Nitride, Synthesis, Neutron Powder Diffraction, Close Packing